

Semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B_{s1}'

Zhi-Gang Luo¹, Xiao-Lin Chen¹, Xiang Liu^{1,2,*} and Shi-Lin Zhu^{1†}

¹*Department of Physics, Peking University, Beijing 100871, China*

²*Centro de Física Teórica, Departamento de Física,
Universidade de Coimbra, P-3004-516 Coimbra, Portugal*

(Dated: February 26, 2009)

Stimulated by recent observations of the excited bottom-strange mesons B_{s1} and B_{s2}^* , we calculate the semileptonic decays B_{s0} , B_{s1}' , B_{s1} , $B_{s2}^* \rightarrow [D_s(1968), D_s^*(2112), D_{sJ}(2317), D_{sJ}(2460)]\ell\bar{\nu}$, which is relevant for the exploration of the potential of searching these semileptonic decays in experiment.

PACS numbers: 12.38.Lg, 12.39.Fe, 12.39.Hg

I. INTRODUCTION

Recently, the CDF collaboration announced the observation of two orbitally excited narrow B_s mesons. Their masses are $m_{B_{s1}} = 5829.4 \pm 0.7$ MeV and $m_{B_{s2}^*} = 5839.6 \pm 0.7$ MeV [1]. Later the D0 collaboration confirmed the B_{s2}^* state with $m_{B_{s2}^*} = 5839.6 \pm 1.1(\text{stat.}) \pm 0.7(\text{syst.})$ MeV [2]. Meanwhile, the D0 collaboration indicated that B_{s1} was not observed with the available data set [2]. In the heavy quark effective theory (HQET), the charmed or bottom mesons can be categorized into doublets since the angular momentum of the light components j_ℓ is a good quantum number in the $m_Q \rightarrow \infty$ limit. They are $j_\ell^P = \frac{1}{2}^-$, the H doublet $(0^-, 1^-)$ with the orbital angular momentum $L = 0$, $j_\ell^P = \frac{1}{2}^+$, the S doublet $(0^+, 1^+)$ and $j_\ell^P = \frac{3}{2}^+$, the T doublet $(1^+, 2^+)$ with $L = 1$. The recently observed B_{s1} and B_{s2}^* mesons belong to the T doublet [1, 2].

The observations of B_{s1} and B_{s2}^* [1, 2] not only enrich the mass spectrum of the bottom-strange system¹ but also inspire our interest in these two unobserved P-wave bottom-strange mesons. Since B_{s1} and B_{s2}^* lie above the thresholds of BK and B^*K , their strong decay is dominant. In our recent work [4], we calculated the strong decays of B_{s1} and B_{s2}^* with the $^3P_0$ model. Our result indicates that the two-body strong decay widths of B_{s1} and B_{s2}^* can reach up to 98 keV and 5 MeV, respectively. In contrast, B_{s0} and B_{s1}' were generally speculated to lie below the threshold of BK and B^*K in Ref. [5]. In fact, they are expected to be narrow resonances with a width around several tens of keV [6] since their main decay modes are the isospin-violating strong decays and electromagnetic decays. These two states are still missing in the experiments. Up to now, they have not been observed experimentally.

The semileptonic decays of B_{s0} , B_{s1}' , B_{s1} , B_{s2}^* have not been explored. In this work, we will calculate the semileptonic decays B_{s0} , B_{s1}' , B_{s1} , $B_{s2}^* \rightarrow [D_s(1968), D_s^*(2112), D_{sJ}(2317), D_{sJ}(2460)]\ell\bar{\nu}$ in the framework of the constituent-quark meson (CQM) model, where $(D_s(1968), D_s^*(2112))$ and $(D_{sJ}(2317), D_{sJ}(2460))$ belong to H and S doublets of the bottom-strange meson respectively.

Through the present investigation, we want to learn the potential of searching the semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B_{s1}' in future experiments such as CDF, D0 and the forthcoming Large Hadron Collider beauty (LHCb). If these semileptonic decays can be reached by the future experiments, one may compare the bottom-strange mesons with these two exotic charm-strange states $D_{sJ}(2317)$ and $D_{sJ}(2460)$ through the semileptonic decays. The observations of the above narrow charm-strange mesons have resulted in an extensive study of their properties in the past five years².

This work is organized as follows. After the introduction, we briefly introduce the theoretical framework, i.e. the CQM model. In Section III, the formulation relevant to the semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B_{s1}' will be

*Electronic address: liuxiang@teor.fis.uc.pt

†Electronic address: zhushl@phy.pku.edu.cn

¹ Besides B_{s1} and B_{s2}^* observed by CDF and D0, there are only two established bottom-strange states (B_s and B_s^*) listed in Particle Data group (PDG) so far [3].

² $D_{sJ}(2317)$ and $D_{sJ}(2460)$ with spin parity structures $J^P = 0^+$ and $J^P = 1^+$ [7, 8, 9] have inspired heated debates about their structures. A detailed review can be found in Ref. [10]. Possible interpretations include the $(0^+, 1^+)$ chiral partners of D_s and D_s^* [11], P-wave excited states of D_s and D_s^* [12], coupled-channel effects between $c\bar{s}$ states and the DK continuum [13], conventional $c\bar{s}$ states [14, 15], four-quark states [17, 18, 19] etc. Considering the large contribution of the S-wave DK continuum in the QCD sum rule (QSR) approach, the mass of $D_{sJ}(2317)$ agrees well with the experimental value [16]. Therefore, $D_{sJ}(2317)$ and $D_{sJ}(2460)$ are very probably conventional $c\bar{s}$ states with $J^P = 0^+$ and $J^P = 1^+$ [10].

given. In Section IV, we present the numerical results with all input parameters. The last section is a short discussion. The detailed expressions are collected in the appendix.

II. THE CQM MODEL

We first give a brief review of the CQM model [20, 21]. The effective Lagrangian of the CQM model incorporates both the heavy-quark spin-flavor symmetry and the chiral symmetry [20]

$$\begin{aligned} \mathcal{L}_{CQM} = & \bar{\chi}[\gamma \cdot (i\partial + \mathcal{V})]\chi + \bar{\chi}\gamma \cdot \mathcal{A}\gamma_5\chi - m_q\bar{\chi}\chi + \frac{f_\pi^2}{8}Tr[\partial^\mu\Sigma\partial_\mu\Sigma^\dagger] + \bar{h}_v(iv \cdot \partial)h_v \\ & - [\bar{\chi}(\bar{H} + \bar{S} + i\bar{T}^\mu\frac{\partial_\mu}{\Lambda})h_v + h.c.] + \frac{1}{2G_3}Tr[(\bar{H} + \bar{S})(H - S)] + \frac{1}{2G_4}Tr[\bar{T}^\mu T_\mu]. \end{aligned} \quad (1)$$

Here H , S and T denote the super-fields corresponding to the $(0^-, 1^-)$, $(0^+, 1^+)$ and $(1^+, 2^+)$ doublets respectively, whose explicit matrix representations are [22]

$$H = \frac{1+\not{v}}{2}[P_\mu^*\gamma^\mu - P\gamma_5], \quad (2)$$

$$S = \frac{1+\not{v}}{2}[P_{1\mu}^*\gamma^\mu\gamma_5 - P_0], \quad (3)$$

$$T^\mu = \frac{1+\not{v}}{2}\left\{P_2^{*\mu\nu}\gamma_\nu - \sqrt{\frac{3}{2}}P_{1\nu}^*\gamma_5[g^{\mu\nu} - \frac{1}{3}\gamma^\nu(\gamma^\mu - v^\mu)]\right\}. \quad (4)$$

P , $P^{*\mu}$, P_0 and $P_1^{*\mu}$ correspond to the annihilation operators of the pseudoscalar, vector, scalar and axial vector mesons respectively. They are normalized as

$$\begin{aligned} \langle 0|P|M(0^-)\rangle &= \sqrt{M_H}, & \langle 0|P^{*\mu}|M(1^-)\rangle &= \sqrt{M_H}\epsilon^\mu, \\ \langle 0|P_0|M(0^+)\rangle &= \sqrt{M_S}, & \langle 0|P_1^{*\mu}|M(1^+)\rangle &= \sqrt{M_S}\epsilon^\mu, \\ \langle 0|P_1^{*\mu}|M(1^+)\rangle &= \sqrt{M_T}\epsilon^\mu, & \langle 0|P_2^{*\mu\nu}|M(2^+)\rangle &= \sqrt{M_T}\eta^{\mu\nu}. \end{aligned}$$

In Eq. (1), the fifth term represents the kinetic term of heavy quarks with $\not{v}h_v = h_v$. $\chi = \xi q$ ($q = u, d, s$) denotes the light-quark field with $\xi = e^{\frac{i\mathcal{M}}{f_\pi}}$ and $\mathcal{M}$ is the octet pseudoscalar matrix. $\mathcal{V}^\mu$ and $\mathcal{A}^\mu$ are defined as

$$\mathcal{V}^\mu = \frac{1}{2}(\xi^\dagger\partial^\mu\xi + \xi\partial^\mu\xi^\dagger), \quad (5)$$

$$\mathcal{A}^\mu = \frac{-i}{2}(\xi^\dagger\partial^\mu\xi - \xi\partial^\mu\xi^\dagger). \quad (6)$$

An important feature of the effective Lagrangian in Eq. (1) is that $\mathcal{L}_{CQM}$ describes the interaction vertex of the heavy-meson with heavy and light quarks, which makes the study of the phenomenology of heavy-meson physics at the quark level feasible. The CQM model has been applied to study the heavy meson phenomenology [23, 24, 25, 26, 27]. In Ref. [26], the semileptonic decays $B_s \rightarrow D_{sJ}(2317, 2460)\ell\bar{\nu}$ have been studied in the CQM model. In this work, we extend the same formalism to the semileptonic decays of B_{s1} and B_{s2}^* . The interested readers may also consult the review paper of the CQM model in [20].

III. THE SEMILEPTONIC DECAYS OF B_{s1} , B_{s2}^* , B_{s0} AND B_{s1}'

In this section, we calculate the semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B_{s1}' in the CQM model. We are interested in the semileptonic decay modes of B_{s1} , B_{s2}^* , B_{s0} and B_{s1}' including $D_s(1968)\ell\bar{\nu}$, $D_s^*(2112)\ell\bar{\nu}$, $D_{sJ}(2317)\ell\bar{\nu}$ and $D_s(2460)\ell\bar{\nu}$, which are depicted in Fig. 1.

The four-fermion operator describing $b \rightarrow c + \ell\bar{\nu}$ is

$$\mathcal{O} = \frac{G_F V_{cb}}{\sqrt{2}}\bar{c}\gamma^\mu(1 - \gamma_5)b\bar{\nu}\gamma_\mu(1 - \gamma_5)\ell. \quad (7)$$




FIG. 1: The Feynman diagram depicting the semileptonic decays of $B_{s1}(B_{s2}^*) \rightarrow D_s \ell \bar{\nu}$. Here the thick-line denotes the heavy-quark propagator.

Thus the general decay amplitudes of $B_{s1}(B_{s2}^*, B_{s0}, B'_{s1}) \rightarrow D_s(D_s^*, D_{sJ}(2317, 2460)) \ell \bar{\nu}$ read

$$\begin{aligned} \mathcal{M}[(b\bar{s}) \rightarrow (c\bar{s}) + \ell\bar{\nu}] \\ = \frac{G_F V_{cb}}{\sqrt{2}} \langle (c\bar{s}) | \bar{c} \gamma^\mu (1 - \gamma_5) b | (b\bar{s}) \rangle \langle \ell\bar{\nu} | \bar{\nu} \gamma_\mu (1 - \gamma_5) \ell | 0 \rangle, \end{aligned} \quad (8)$$

where we use $(c\bar{s})$ and $(b\bar{s})$ to denote the charm-strange and bottom-strange mesons, respectively. The hadronic matrix element $\langle (c\bar{s}) | \bar{c} \gamma^\mu (1 - \gamma_5) b | (b\bar{s}) \rangle$ has to be calculated using phenomenological models, such as QSR and the CQM model. Here we adopt the CQM model to calculate the hadronic matrix element. Figure 1 illustrates the semileptonic decays $B_{s1}(B_{s2}^*) \rightarrow D_s \ell \bar{\nu}$ in the CQM model.

Interaction	Vertex
$b - B_{s1} - \bar{s}$	$(\frac{i}{\Lambda}) \sqrt{\frac{3}{2}} \sqrt{m_{B_{s1}} Z_T} \frac{1+\not{v}}{2} \gamma_5 [\epsilon \cdot k - \frac{\not{\epsilon}}{3} (\not{k} - v \cdot k)]$
$b - B_{s2}^* - \bar{s}$	$(-\frac{i}{\Lambda}) \sqrt{m_{B_{s2}^*} Z_T} \frac{1+\not{v}}{2} k_\mu \eta^{\mu\nu} \gamma_\nu$
$b - B_{s0} - \bar{s}$	$i \sqrt{m_{B_{s0}}^* Z_S} \frac{1+\not{v}}{2}$
$b - B'_{s1} - \bar{s}$	$-i \sqrt{m_{B'_{s1}}^* Z_S} \not{\epsilon}_2 \gamma_5 \frac{1+\not{v}}{2}$
$c - D_s - \bar{s}$	$-i \sqrt{m_{D_s} Z_H} \gamma_5 \frac{1+\not{v}'}{2}$
$c - D_s^* - \bar{s}$	$-i \sqrt{m_{D_s} Z_H} \not{\epsilon}_1^* \frac{1+\not{v}'}{2}$
$c - D_{s0} - \bar{s}$	$i \sqrt{m_{D_{s0}}^* Z_S} \frac{1+\not{v}'}{2}$
$c - D'_{s1} - \bar{s}$	$-i \sqrt{m_{D'_{s1}}^* Z_S} \not{\epsilon}_2^* \gamma_5 \frac{1+\not{v}'}{2}$

TABLE I: The interaction vertex for the semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B'_{s1} .

According to Eq. (1), we give the interaction vertices in Table I, which are related to the semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B'_{s1} . Here ϵ , η , ϵ_1 and ϵ_2 denote the polarization vectors of the heavy mesons. Note that we use D_{s0} and D'_{s1} to denote $D_{sJ}(2317)$ and $D_{sJ}(2460)$ respectively. The normalization constants $Z_{H,S,T}$ are given in Ref. [20]:

$$Z_H^{-1} = (\Delta_H + m_s) \frac{\partial \mathcal{I}_3(\Delta_H)}{\partial \Delta_H} + \mathcal{I}_3(\Delta_H), \quad (9)$$

$$Z_S^{-1} = (\Delta_S - m_s) \frac{\partial \mathcal{I}_3(\Delta_S)}{\partial \Delta_S} + \mathcal{I}_3(\Delta_S), \quad (10)$$

$$\begin{aligned} Z_T^{-1} = \frac{1}{3\Lambda^2} \Big\{ (\Delta_T^2 - m_s^2) [\mathcal{I}_3(\Delta_T) + (m_s + \Delta_T) \frac{\partial \mathcal{I}_3(\Delta_T)}{\partial \Delta_T}] \\ + (m_s + \Delta_T) [\mathcal{I}_1 + 2\Delta_T \mathcal{I}_3(\Delta_T) + \frac{\partial \mathcal{I}_0(\Delta_T)}{\partial \Delta_T}] + \mathcal{I}_0(\Delta_T) + \Delta_T \mathcal{I}_1 \Big\} \end{aligned} \quad (11)$$

with the definitions of $I_{0,1,3}$ listed in the appendix.

In terms of the above expressions, we can write the general expression of the hadronic matrix element $\langle(c\bar{s})|\bar{c}\gamma^\mu(1-\gamma_5)b|(b\bar{s})\rangle$ in the CQM model

$$\begin{aligned} & \langle(c\bar{s})|\bar{c}\gamma^\mu(1-\gamma_5)b|(b\bar{s})\rangle \\ &= -\frac{iN_c}{16\pi^4} \int d^4l \frac{Tr[\gamma^\mu(1-\gamma_5)\Gamma_a(\gamma \cdot l + m_s)\Gamma_b]}{(l^2 - m_s^2)(v \cdot l + \alpha)(v' \cdot l + \beta)}, \end{aligned} \quad (12)$$

where Γ_a corresponds to the vertex of the interaction of $B_{s1}(B_{s2}^*, B_{s0}, B'_{s1})$ with b and $\bar{s}$. Γ_b is the vertex describing the interaction of charm-strange meson with c and $\bar{s}$ quarks. N_c denotes the colors degrees of freedom and $N_c = 3$. $\alpha(\beta) = \Delta_{H,S,T}$ denotes the mass difference between the heavy mesons and the heavy quark [20].

In the following, by substituting Γ_a and Γ_b with the expression listed in Table I, one obtains the hadron matrix elements relevant to the semileptonic decays of B_{s1} and B_{s2}^* with the transitions of $B_{s1}(B_{s2}^*) \rightarrow D_{s0}(D'_{s1})$:

$$\begin{aligned} & \langle D_{s0}(v')|\gamma^\mu(1-\gamma_5)|B_{s1}(v, \epsilon)\rangle \\ &= \sqrt{m_{B_{s1}}m_{D_{s0}}}\zeta(\omega) \left[f\epsilon^\mu + (\epsilon \cdot v')(h_1v^\mu + h_2v'^\mu) + ih_3\varepsilon^{\mu\delta\lambda\rho}v'_\delta v_\lambda \epsilon_\rho \right], \end{aligned} \quad (13)$$

$$\begin{aligned} & \langle D'_{s1}(v', \epsilon_2)|\gamma^\mu(1-\gamma_5)|B_{s1}(v, \epsilon)\rangle \\ &= \sqrt{m_{B_{s1}}m_{D'_{s1}}}\zeta(\omega) \left[f_1(v \cdot \epsilon_2^*)\epsilon^\mu + f_2(v' \cdot \epsilon)\epsilon_2^{*\mu} + h_1(\epsilon_2^* \cdot \epsilon)v^\mu + h_2(\epsilon_2^* \cdot \epsilon)v'^\mu + h'_2(v' \cdot \epsilon)(v \cdot \epsilon_2^*)v'^\mu \right. \\ & \quad \left. + ih_3\varepsilon^{\mu\delta\lambda\rho}\epsilon_{2\delta}^*v_\lambda \epsilon_\rho + ih_4\varepsilon^{\mu\delta\lambda\rho}\epsilon_{2\delta}^*v'_\lambda \epsilon_\rho + ih(v' \cdot \epsilon)\varepsilon^{\mu\delta\lambda\rho}\epsilon_{2\delta}^*v'_\lambda v_\rho \right] \end{aligned} \quad (14)$$

for the semileptonic decays of B_{s1} ,

$$\begin{aligned} & \langle D_{s0}(v')|\gamma^\mu(1-\gamma_5)|B_{s2}^*(v, \eta)\rangle \\ &= \sqrt{m_{B_{s2}^*}m_{D_{s0}}}\zeta(\omega) \left[f v'^\alpha g^{\beta\mu} + g v'^\alpha v^\mu v'^\beta + ih\varepsilon^{\beta\mu\delta\lambda}v_\delta v'_\lambda v'^\alpha \right] \eta_{\alpha\beta}, \end{aligned} \quad (15)$$

$$\begin{aligned} & \langle D'_{s1}(v', \epsilon_2)|\gamma^\mu(1-\gamma_5)|B_{s2}^*(v, \eta)\rangle \\ &= \sqrt{m_{B_{s2}^*}m_{D'_{s1}}}\zeta(\omega) \left[f v'^\alpha v'^\beta \epsilon_2^{*\mu} + f_1 v'^\alpha \epsilon_2^{*\beta} v^\mu + f_2 v'^\alpha \epsilon_2^{*\beta} v'^\mu \right. \\ & \quad \left. + g g^{\beta\mu}(\epsilon_2^* \cdot v)v'^\alpha + ih_1\varepsilon^{\beta\mu\delta\lambda}\epsilon_{2\delta}^*v_\lambda v'^\alpha + ih_2\varepsilon^{\beta\mu\delta\lambda}\epsilon_{2\delta}^*v'_\lambda v'^\alpha \right] \eta_{\alpha\beta} \end{aligned} \quad (16)$$

for the semileptonic decays of B_{s2}^* . Here $\omega = v \cdot v'$.

$B_{s1} \rightarrow$	$\sqrt{6}f$	$\sqrt{6}f_1$	$\sqrt{6}f_2$	$\sqrt{6}h_1$	$\sqrt{6}h_2$	$\sqrt{6}h'_2$	$\sqrt{6}h_3$	$\sqrt{6}h_4$	$\sqrt{6}h$
D_s	$-(\omega^2 - 1)$	-	-	$\omega - 2$	-3	-	$-(\omega + 1)$	-	-
D_{s0}	$-(\omega^2 - 1)$	-	-	$\omega + 2$	-3	-	$-(\omega - 1)$	-	-
D_s^*	-	$\omega + 1$	$2(\omega + 1)$	$-(\omega + 1)$	$\omega + 1$	-3	$\omega + 1$	$-(\omega + 1)$	3
D'_{s1}	-	$-(\omega - 1)$	$2(\omega - 1)$	$\omega - 1$	$\omega - 1$	-3	$-(\omega - 1)$	$-(\omega - 1)$	3

TABLE II: The coefficients relevant to the transitions of $B_{s1} \rightarrow D_s, D_s^*, D_{s0}, D'_{s1}$.

It is worth noting that the Lorentz structures of the matrix elements $B_{s1}(B_{s2}^*) \rightarrow D_s$ and $B_{s1}(B_{s2}^*) \rightarrow D_s^*$ are similar to those of the transitions of $B_{s1}(B_{s2}^*) \rightarrow D_{s0}$ and $B_{s1}(B_{s2}^*) \rightarrow D'_{s1}$ respectively. For different semileptonic decay, the coefficients $f_i, h_i^{(\prime)}, g, h$ are different; they are given in Tables II-III. Meanwhile, the form factor $\zeta(\omega)$ for

$B_{s2}^* \rightarrow$	f	f_1	f_2	g	h	h_1	h_2
D_s	$\omega + 1$	-	-	-1	1	-	-
D_{s0}	$\omega - 1$	-	-	-1	1	-	-
D_s^*	-1	1	1	-1	-	1	1
D'_{s1}	-1	-1	1	1	-	-1	1

TABLE III: The coefficients relevant to the transitions of $B_{s2}^* \rightarrow D_s, D_s^*, D_{s0}, D'_{s1}$.

$B_{s1}(B_{s2}^*) \rightarrow D_{s0}(D'_{s1})$ should be replaced by $\xi(\omega)$ for $B_{s1}(B_{s2}^*) \rightarrow D_s(D_s^*)$. In the above expressions of the hadron element matrices, there exist two independent form factors $\xi(\omega)$ and $\zeta(\omega)$:

$$\xi(\omega) = \Lambda^{-1} \sqrt{Z_T Z_H} [\mathcal{A}_2 + \mathcal{A}_3 - \mathcal{B}_2 m_s]_{\alpha=\Delta_T, \beta=\Delta_H}, \quad (17)$$

$$\zeta(\omega) = \Lambda^{-1} \sqrt{Z_T Z_S} [\mathcal{A}_2 - \mathcal{A}_3 + \mathcal{B}_2 m_s]_{\alpha=\Delta_T, \beta=\Delta_S}. \quad (18)$$

Equations (13)-(16) are consistent with the heavy-quark spin-flavor symmetry. The expressions of $\mathcal{A}_2$, $\mathcal{A}_3$ and $\mathcal{B}_2$ are given in the appendix. Finally, the differential rates of the semileptonic decays of B_{s1} and B_{s2}^* are

$$\frac{d\Gamma}{d\omega}(B_{sJ} \rightarrow D_{sJ'} \ell \bar{\nu}) = \frac{G_F^2 |V_{cb}|^2}{(2J+1)48\pi^3} m_{B_{sJ}} r^2 (\omega^2 - 1)^{1/2} \Theta[B_{sJ}, D_{sJ'}] \quad (19)$$

with

$$\Theta[B_{sJ}, D_{sJ'}] = (q_\mu q_\nu - q^2 g_{\mu\nu}) W_J^{\mu\nu}(D_{sJ'}), \quad (20)$$

where

$$W_J^{\mu\nu}(D_{sJ'}) = \sum_{spins} \langle D_{sJ'} | \gamma^\mu (1 - \gamma_5) | B_{sJ} \rangle \langle D_{sJ'} | \gamma^\nu (1 - \gamma_5) | B_{sJ} \rangle^*, \quad (21)$$

and $r = m_{D_{sJ'}}/m_{B_{sJ}}$. The allowed integral range for ω is

$$0 \leq \omega - 1 \leq \frac{(m_{B_{sJ}} - m_{D_{sJ'}})^2}{2m_{B_{sJ}} m_{D_{sJ'}}}. \quad (22)$$

For the processes of $B_{s1} \rightarrow D_s \ell \bar{\nu}$, $B_{s1} \rightarrow D_s^* \ell \bar{\nu}$, $B_{s1} \rightarrow D_{s0} \ell \bar{\nu}$ and $B_{s1} \rightarrow D'_{s1} \ell \bar{\nu}$, the expressions of $\Theta[B_{s1}, D_{sJ'}]$ are respectively

$$\begin{aligned} & \Theta[B_{s1}, D_{s0}(D_s)] \\ &= m_{B_{s1}}^4 r |\zeta(\omega)|^2 \left\{ [(r\omega - 1)f + (\omega^2 - 1)(rh_1 + h_2)]^2 + 2(r^2 - 2r\omega + 1)[f^2 + (\omega^2 - 1)h_3^2] \right\}, \end{aligned} \quad (23)$$

$$\begin{aligned} & \Theta[B_{s1}, D'_{s1}(D_s^*)] \\ &= m_{B_{s1}}^4 r |\zeta(\omega)|^2 \left\{ (\omega^2 - 1) \{ [(r\omega - 1)f_1 - (r - \omega)f_2 + \omega(rh_1 + h_2) + (\omega^2 - 1)h_2']^2 + 2(rh_1 + h_2)^2 \right. \\ & \quad + 2(1 + r^2 - 2r\omega)(f_1^2 + f_2^2) \} + 2[(r\omega - 1)h_3 + (r - \omega)h_4]^2 + 2(1 + r^2 - 2r\omega) \{ (\omega h_3 + h_4)^2 \\ & \quad \left. + [h_3 + \omega h_4 + (\omega^2 - 1)h]^2 \} \right\}. \end{aligned} \quad (24)$$

For the semileptonic decays of B_{s2}^* , the $\Theta[B_{s1}, D_{sJ'}]$ read

$$\begin{aligned} & \Theta[B_{s2}^*, D_{s0}(D_s)] \\ &= m_{B_{s2}}^4 r |\zeta(\omega)|^2 (\omega^2 - 1) \left\{ \frac{2}{3} [(r\omega - 1)f + r(\omega^2 - 1)g]^2 + (1 + r^2 - 2r\omega)[f^2 + (\omega^2 - 1)h^2] \right\}, \end{aligned} \quad (25)$$

$$\begin{aligned}
& \Theta[B_{s2}^*, D'_{s1}(D_s^*)] \\
& = m_{B_{s2}}^4 r |\zeta(\omega)|^2 (\omega^2 - 1) \left\{ (\omega^2 - 1) \left\{ \frac{2}{3} [-(r - \omega)f + \omega(rf_1 + f_2) + (r\omega - 1)g]^2 + (rf_1 + f_2)^2 \right. \right. \\
& \quad \left. \left. + (rh_1 + h_2)^2 + (1 + r^2 - 2r\omega) \left(\frac{4}{3} f^2 + g^2 + h_1^2 - 2h_2^2 \right) \right\} + \frac{10}{3} (1 + r^2 - 2r\omega) (h_1 + \omega h_2)^2 \right\}. \quad (26)
\end{aligned}$$

$B_{s0} \rightarrow$	f	h_1	h_2	h_3
D_s	-	1	-1	-
D_{s0}	-	-1	-1	-
D_s^*	$1 - \omega$	0	1	1
D'_{s1}	$\omega + 1$	0	-1	1

TABLE IV: The coefficients relevant to the transitions of $B_{s0} \rightarrow D_s, D_s^*, D_{s0}, D'_{s1}$.

$B'_{s1} \rightarrow$	f	f_1	f_2	h_1	h_2	h'_2	h_3	h_4	h
D_s	$\omega - 1$	-	-	-1	0	-	1	-	-
D_{s0}	$\omega + 1$	-	-	1	0	-	1	-	-
D_s^*	-	-1	1	1	-1	0	-1	1	0
D'_{s1}	-	-1	-1	1	1	0	-1	-1	0

TABLE V: The coefficients relevant to the transitions of $B'_{s1} \rightarrow D_s, D_s^*, D_{s0}, D'_{s1}$.

In this work, we also calculate the semileptonic decays of the B_{s0} and B'_{s1} mesons in the S doublet, i.e. $B_{s0}(B'_{s1}) \rightarrow D_s(D_s^*, D_{s0}, D'_{s1})\ell\bar{\nu}$. For $B_{s0} \rightarrow D_{s0}(D'_{s1})\ell\bar{\nu}$, we have

$$\Theta[B_{s0}, D_{s0}] = m_{B_{s0}}^4 r |\chi(\omega)|^2 (\omega^2 - 1) (rh_1 + h_2)^2, \quad (27)$$

$$\begin{aligned}
& \Theta[B_{s0}, D'_{s1}] \\
& = m_{B_{s0}}^4 r |\chi(\omega)|^2 \{ [(\omega - r)f + (\omega^2 - 1)(rh_1 + h_2)]^2 + 2(r^2 - 2r\omega + 1)[f^2 + (\omega^2 - 1)h_3^2] \}, \quad (28)
\end{aligned}$$

where

$$\chi(\omega) = Z_S [\mathcal{B}_1 + \mathcal{B}_2 + m_s]_{\alpha=\Delta_S, \beta=\Delta_S}. \quad (29)$$

The $\Theta[B'_{s1}, D_{s0}]$ and $\Theta[B'_{s1}, D'_{s1}]$ functions for the $B'_{s1} \rightarrow D_{s0}(D'_{s1})\ell\bar{\nu}$ decays are similar to Eqs. (23)-(24) where $\zeta(\omega)$ has to be replaced by the new form factor $\chi(\omega)$. The corresponding parameters and coefficients are listed in Tables IV-V.

For $B_{s0} \rightarrow D_s(D_s^*)\ell\bar{\nu}$, $\Theta[B_{s0}, D_s]$ and $\Theta[B_{s0}, D_s^*]$ can be obtained after replacing $\chi(\omega)$ by $\lambda(\omega)$ in Eq. (27)-(28), where

$$\lambda(\omega) = \sqrt{Z_H Z_S} [\mathcal{B}_1 - \mathcal{B}_2 + m_s]_{\alpha=\Delta_S, \beta=\Delta_H}. \quad (30)$$

Similarly, one gets the functions $\Theta[B'_{s1}, D_s]$ and $\Theta[B'_{s1}, D_s^*]$ for $B'_{s1} \rightarrow D_s(D_s^*)\ell\bar{\nu}$ with the replacement $\zeta(\omega) \rightarrow \lambda(\omega)$ in Eqs. (23)-(24).

IV. NUMERICAL RESULTS

We now collect the input parameters: $G_F = 1.1664 \times 10^{-5} \text{ GeV}^{-2}$, $V_{cb} = 0.043$; $M_{B_{s1}} = 5829.4 \text{ MeV}$, $M_{B_{s2}^*} = 5839.6 \text{ MeV}$, $M_{D_s} = 1968.2 \text{ MeV}$, $M_{D_s^*} = 2112.0 \text{ MeV}$, $M_{D_{sJ}^*(2317)} = 2317.3 \text{ MeV}$, $M_{D_{sJ}^*(2460)} = 2458.9 \text{ MeV}$ [3]. $M_{B_{s0}} = 5718 \text{ MeV}$ and $M_{B'_{s1}} = 5765$ [5]. $m_s = 0.5 \text{ GeV}$, $\Lambda = 1.25 \text{ GeV}$, the infrared cutoff $\mu = 0.593 \text{ GeV}$ and $\Delta_S - \Delta_H = 335 \pm 35 \text{ MeV}$ [27].

In Table VI, we give the ranges of $\Delta_{H,S,T}$ for the strange sector, which are given in Ref. [27, 28]. In terms of the definitions of $Z_{H,S,T}$ in Eqs. (9)-(11), we obtain the values of $Z_{H,S,T}$ listed in Table VI.

Type	Δ_H	Δ_S	Δ_T	Z_H	Z_S	Z_T
(a)	0.5	0.86	0.84	4.87	2.95	3.26
(b)	0.6	0.91	0.94	3.45	2.28	1.91
(c)	0.7	0.97	1.04	2.37	1.66	1.06

TABLE VI: The values of $\Delta_{H,S,T}$ and the corresponding $Z_{H,S,T}$. Here $\Delta_{H,S,T}$ and $Z_{H,S,T}$ are in units of GeV and GeV^{-1} , respectively.




FIG. 2: (a) The dependence of $\xi(\omega)$ on ω . Here the solid, dashed and dotted lines correspond to the results with parameters $(\Delta_T = 0.84, \Delta_H = 0.5)$, $(\Delta_T = 0.94, \Delta_H = 0.6)$, $(\Delta_T = 1.04, \Delta_H = 0.7)$, respectively. (b) The variation of $\zeta(\omega)$ with ω . Here the solid, dashed and dotted lines correspond to the results with parameters $(\Delta_T = 0.84, \Delta_S = 0.86)$, $(\Delta_T = 0.94, \Delta_S = 0.91)$, $(\Delta_T = 1.04, \Delta_S = 0.97)$, respectively. (c) The dependence of $\chi(\omega)$ on ω . Here the solid, dashed and dotted lines correspond to the results with parameters $\Delta_S = 0.86$, $\Delta_S = 0.91$, $\Delta_S = 0.97$, respectively. (d) The variation of $\lambda(\omega)$ with ω . Here the solid, dashed and dotted lines correspond to the results with parameters $(\Delta_S = 0.86, \Delta_H = 0.5)$, $(\Delta_S = 0.91, \Delta_H = 0.6)$, $(\Delta_S = 0.97, \Delta_H = 0.7)$, respectively.

With the above parameters, Fig. 2 illustrates the dependence of the form factors $\xi(\omega)$, $\zeta(\omega)$, $\chi(\omega)$ and $\lambda(\omega)$ on ω .

In Tables VII and VIII, we give the decay widths of the semileptonic decays of B_{s1} , B_{s2}^* , B_{s0} and B_{s1}' with the three different choices of parameters $\Delta_{H,S,T}$ listed in Table VI. Up to now, the CDF and D0 collaborations have not measured the decay widths of B_{s1} and B_{s2}^* .

Using the semileptonic decays of B_{s1} as an example, we show the variation of the differential rates of the semileptonic decays of B_{s1} with ω in Fig. 3. The differential rates of $B_{s1} \rightarrow D_s \ell \bar{\nu}$ and $B_{s1} \rightarrow D_s^* \ell \bar{\nu}$ increase with ω monotonically.

	$B_{s1} \rightarrow$			$B_{s2}^* \rightarrow$		
	(a)	(b)	(c)	(a)	(b)	(c)
$D_s \ell \bar{\nu}$	2.1×10^{-15}	1.8×10^{-15}	1.6×10^{-15}	2.1×10^{-15}	1.8×10^{-15}	1.6×10^{-15}
$D_s^* \ell \bar{\nu}$	4.9×10^{-15}	4.4×10^{-15}	3.9×10^{-15}	5.0×10^{-15}	4.5×10^{-15}	3.9×10^{-15}
$D_{s0} \ell \bar{\nu}$	8.7×10^{-20}	4.7×10^{-20}	6.9×10^{-20}	5.6×10^{-20}	3.0×10^{-20}	4.5×10^{-20}
$D'_{s1} \ell \bar{\nu}$	1.0×10^{-19}	8.7×10^{-20}	1.5×10^{-19}	1.2×10^{-19}	1.0×10^{-19}	1.8×10^{-19}

TABLE VII: The decay widths of the semileptonic decays of B_{s1} and B_{s2}^* . Here columns (a), (b), (c) correspond to the results with the three parameter combinations listed in Table VI. The decay widths are in units of GeV.

	$B_{s0} \rightarrow$			$B'_{s1} \rightarrow$		
	(a)	(b)	(c)	(a)	(b)	(c)
$D_s \ell \bar{\nu}$	2.5×10^{-14}	1.3×10^{-14}	5.9×10^{-15}	1.2×10^{-14}	6.4×10^{-15}	2.9×10^{-15}
$D_s^* \ell \bar{\nu}$	2.5×10^{-15}	1.3×10^{-14}	5.9×10^{-15}	3.8×10^{-14}	2.0×10^{-14}	9.1×10^{-15}
$D_{s0} \ell \bar{\nu}$	1.6×10^{-15}	3.9×10^{-16}	2.8×10^{-15}	4.9×10^{-16}	9.6×10^{-16}	3.7×10^{-15}
$D'_{s1} \ell \bar{\nu}$	1.5×10^{-15}	3.3×10^{-15}	1.3×10^{-14}	2.1×10^{-15}	2.7×10^{-15}	1.2×10^{-14}

TABLE VIII: The decay widths of the semileptonic decays of B_{s0} and B'_{s1} . Here columns (a), (b), (c) correspond to the results with the three parameter combinations listed in Table VI. The decay widths are in units of GeV.

In contrast, the variation of the differential rates of $B_{s1} \rightarrow D_{s0} \ell \bar{\nu}$ and $B_{s1} \rightarrow D'_{s1} \ell \bar{\nu}$ with ω are not monotonic. The different line shapes of the differential rates of the semileptonic decays of B_{s1} account for the results of the decay widths in Table VII, i.e. it tells us why the decay rates for type (c) are not always larger than the results for the other types. The same observation holds for the semileptonic decays of B_{s2}^* , B_{s0} , B'_{s1} .




FIG. 3: (a), (b), (c) and (d) illustrate the dependence of the differential rates of the semileptonic decays of B_{s1} on ω . Here the solid, dashed and dotted lines correspond to type (a), (b) and (c) in Table VI.

V. DISCUSSION

The semileptonic decay of the bottom-strange meson is an interesting topic. Due to the lack of experimental information of the excited bottom-strange mesons, theorists mainly focused on the semileptonic decay of B_s using theoretical approaches such as the QCD sum rule approach [29] and the quark model [26]. The observations of B_{s1} and B_{s2}^* [1, 2] enrich the mass spectrum of the bottom-strange system greatly. In this work, in order to explore the possibility of searching the semileptonic decays of the excited bottom-strange states in the experiments, we have calculated B_{s0} , B'_{s1} , B_{s1} , $B_{s2}^* \rightarrow [D_s(1968), D_s^*(2112), D_{sJ}(2317), D_{sJ}(2460)]\ell\bar{\nu}$ in the framework of the CQM model.

Our numerical results indicate that (1) the decay width of $B_{s1}(B_{s2}^*) \rightarrow D_s(D_s^*)\ell\bar{\nu}$ is around 10^{-15} GeV, which is 4 $\sim$ 5 orders of magnitude larger than that of $B_{s1}(B_{s2}^*) \rightarrow D_{sJ}(2317, 2460)\ell\bar{\nu}$; (2) the decay width of $B_{s0}(B'_{s1}) \rightarrow D_s(D_s^*, D_{sJ}(2317, 2460))\ell\bar{\nu}$ is $10^{-14} \sim 10^{-15}$ GeV.

Although the CDF and D0 experiments did not measure the decay widths of B_{s1} and B_{s2}^* and the experiments did not observe B_{s0} and B'_{s1} , we can roughly estimate the branching ratio of the semileptonic decay of B_{s1} , B_{s2} , B_{s0} and B'_{s1} . In Ref. [4], one obtains the two-body strong decay widths of B_{s1} and B_{s2}^* as 98 keV and 5 MeV respectively. B_{s0} and B'_{s1} are expected to be narrow resonances with a width around several tens of keV³ [6], since their main decay modes are the isospin violating strong decays and electromagnetic decays. Thus it is reasonable to take the strong decay width as the total width approximately for B_{s1} , B_{s2} , B_{s0} and B'_{s1} . We further show the order of magnitude of the semileptonic decay of B_{s1} , B_{s2} , B_{s0} and B'_{s1} in Table IX, which is convenient for the experimentalist to conclude whether the current and future experiments can reach these semileptonic decays.

	$B_{s0} \rightarrow$	$B'_{s1} \rightarrow$	$B_{s1} \rightarrow$	$B_{s2}^* \rightarrow$
$D_s\ell\bar{\nu}$	$10^{-9} \sim 10^{-10}$	$10^{-9} \sim 10^{-10}$	$\sim 10^{-11}$	$\sim 10^{-13}$
$D_s^*\ell\bar{\nu}$	$10^{-9} \sim 10^{-10}$	$10^{-9} \sim 10^{-10}$	$\sim 10^{-11}$	$\sim 10^{-13}$
$D_{s0}\ell\bar{\nu}$	$10^{-10} \sim 10^{-11}$	$10^{-10} \sim 10^{-11}$	$\sim 10^{-16}$	$\sim 10^{-18}$
$D'_{s1}\ell\bar{\nu}$	$10^{-9} \sim 10^{-10}$	$10^{-9} \sim 10^{-10}$	$\sim 10^{-16}$	$\sim 10^{-17}$

TABLE IX: The estimation of the branching fractions of the semileptonic decays of B_{s0} , B'_{s1} , B_{s1} and B_{s2}^* according to our numerical result shown in Table VII and VIII.

From Table IX, we can exclude the possibility of finding the semileptonic decay of $B_{s2}^* \rightarrow [D_s, D_s^*, D_{s0}, D'_{s1}]\ell\bar{\nu}$ and $B_{s1} \rightarrow [D_{s0}, D'_{s1}]\ell\bar{\nu}$ in experiments. However, for $B_{s1} \rightarrow [D_s, D_s^*]\ell\bar{\nu}$ and the semileptonic decays of B_{s0} and B'_{s1} , the upper limit of the branching ratio can reach to 10^{-9} . The present precision of the experimental measurement of the branching fraction of the B mesons has reached up to $10^{-7} \sim 10^{-8}$ [3]. The decays $B_{s1} \rightarrow [D_s, D_s^*]\ell\bar{\nu}$ and the semileptonic decays of B_{s0} and B'_{s1} may be observed in future experiments. Especially, the forthcoming LHCb experiments will produce an enormous amount of data of heavy-flavor hadrons, which is one of the potential experiments in which to search the $B_{s1} \rightarrow [D_s, D_s^*]\ell\bar{\nu}$ and the semileptonic decays of B_{s0} and B'_{s1} . Then these semileptonic decays will be helpful to further test the structure of $D_{sJ}(2317)$ and $D_{sJ}(2460)$. In our calculation, we have assumed $D_{sJ}(2317)$ and $D_{sJ}(2460)$ as the charm-strange mesons with $J^P = 0^+$ and 1^+ . If the experimental measurement of these semileptonic decays are consistent with our prediction, it will provide strong support of the $c\bar{s}$ structure for $D_{sJ}(2317)$ and $D_{sJ}(2460)$.

Acknowledgments

This project was supported by the National Natural Science Foundation of China under Grants 10625521, 10721063, 10705001. X.L. was also supported by the *Fundação para a Ciência e a Tecnologia of the Ministério da Ciência, Tecnologia e Ensino Superior* of Portugal (SFRH/BPD/34819/2007).

³ In Ref. [6], authors calculated the isospin-violating strong decay widths of B_{s0} and B'_{s1} , which are 35 keV and 38 keV, respectively.

Appendix

The definitions of $\mathcal{I}_0(\alpha)$, $\mathcal{I}_1$, $\mathcal{I}_5(\alpha, \beta, \omega)$, $\mathcal{I}_6(\alpha, \beta, \omega)$, $\mathcal{I}_3(\alpha)$, $\mathcal{A}_2(\alpha, \beta, \omega)$, $\mathcal{A}_3(\alpha, \beta, \omega)$, $\mathcal{B}_1(\alpha, \beta, \omega)$ and $\mathcal{B}_2(\alpha, \beta, \omega)$ are [20]

$$\begin{aligned} \mathcal{A}_2(\alpha, \beta, \omega) &= [(\omega^2 - 1)T(\alpha, \beta, \omega) - 6\omega U(\alpha, \beta, \omega) \\ &\quad + (2\omega^2 + 1)S(\alpha, \beta, \omega) + 3S(\beta, \alpha, \omega)]/[2(\omega^2 - 1)^2], \end{aligned} \quad (31)$$

$$\begin{aligned} \mathcal{A}_3(\alpha, \beta, \omega) &= [-\omega(\omega^2 - 1)T(\alpha, \beta, \omega) + 2(2\omega^2 + 1)U(\alpha, \beta, \omega) \\ &\quad - 3\omega(S(\alpha, \beta, \omega) + S(\beta, \alpha, \omega))]/[2(\omega^2 - 1)^2] \end{aligned} \quad (32)$$

$$\mathcal{B}_1(\alpha, \beta, \omega) = [\omega X(\beta, \alpha, \omega) - X(\alpha, \beta, \omega)]/[\omega^2 - 1], \quad (33)$$

$$\mathcal{B}_2(\alpha, \beta, \omega) = [\omega X(\alpha, \beta, \omega) - X(\beta, \alpha, \omega)]/[\omega^2 - 1], \quad (34)$$

with

$$T(\alpha, \beta, \omega) = \mathcal{I}_6(\alpha, \beta, \omega) + m_s^2 \mathcal{I}_5(\alpha, \beta, \omega), \quad (35)$$

$$U(\alpha, \beta, \omega) = \mathcal{I}_1 + \alpha \mathcal{I}_3(\alpha) + \beta \mathcal{I}_3(\beta) + \alpha \beta \mathcal{I}_5(\alpha, \beta, \omega), \quad (36)$$

$$S(\alpha, \beta, \omega) = \omega[\mathcal{I}_1 + \beta \mathcal{I}_3(\beta)] + \alpha \mathcal{I}_3(\beta) + \alpha^2 \mathcal{I}_5(\alpha, \beta, \omega), \quad (37)$$

$$X(\alpha, \beta, \omega) = -\mathcal{I}_3(\beta) - \alpha \mathcal{I}_5(\alpha, \beta, \omega), \quad (38)$$

$$\begin{aligned} \mathcal{I}_0(\alpha) &= \frac{N_c}{16\pi^{3/2}} \int_{1/\Lambda^2}^{1/\mu^2} \frac{dy}{y^{3/2}} e^{-y(m_s^2 - \alpha^2)} \left(\frac{3}{2y} + m_s^2 - \alpha^2 \right) \\ &\quad \times [1 + \text{erf}(\alpha\sqrt{y})] - \alpha \frac{N_c m_s^2}{16\pi^2} \Gamma\left(-1, \frac{m_s^2}{\Lambda^2}, \frac{m_s^2}{\mu^2}\right), \end{aligned} \quad (39)$$

$$\mathcal{I}_1 = \frac{N_c m_s^2}{16\pi^2} \Gamma\left(-1, \frac{m_s^2}{\Lambda^2}, \frac{m_s^2}{\mu^2}\right), \quad (40)$$

$$\mathcal{I}_3(\alpha) = \frac{N_c}{16\pi^{3/2}} \int_{1/\Lambda^2}^{1/\mu^2} \frac{dy}{y^{3/2}} \exp[-y(m_s^2 - \alpha^2)] (1 + \text{erf}(\alpha\sqrt{y})), \quad (41)$$

$$\begin{aligned}
& \mathcal{I}_5(\alpha, \beta, \omega) \\
&= \int_0^1 dx \frac{1}{1 + 2x^2(1 - \omega) + 2x(\omega - 1)} \left[\frac{2N_c}{16\pi^{3/2}} \int_{1/\Lambda^2}^{1/\mu^2} dy \sigma e^{-y(m_s^2 - \sigma^2)} y^{-1/2} (1 + \operatorname{erf}(\sigma\sqrt{y})) \right. \\
&\quad \left. + \frac{2N_c}{16\pi^2} \int_{1/\Lambda^2}^{1/\mu^2} dy e^{-ym_s^2} y^{-1} \right],
\end{aligned} \tag{42}$$

$$\begin{aligned}
& \mathcal{I}_6(\alpha, \beta, \omega) \\
&= \mathcal{I}_1 \int_0^1 dx \frac{\sigma}{1 + 2x^2(1 - \omega) + 2x(\omega - 1)} - \frac{N_c}{16\pi^{3/2}} \int_0^1 dx \frac{1}{1 + 2x^2(1 - \omega) + 2x(\omega - 1)} \\
&\quad \times \int_{1/\Lambda^2}^{1/\mu^2} dy y^{-3/2} e^{-y(m_s^2 - \sigma^2)} \left\{ \sigma [1 + \operatorname{erf}(\sigma\sqrt{y})] [1 + 2y(m_s^2 - \sigma^2)] + 2\sqrt{\frac{y}{\pi}} \left[\frac{3}{2y} + (m_s^2 - \sigma^2) \right] \right\},
\end{aligned} \tag{43}$$

where

$$\sigma(\alpha, \beta, \omega) = \frac{(1 - x)\alpha + x\beta}{\sqrt{1 + 2x^2(1 - \omega) + 2x(\omega - 1)}}.$$

-
- [1] CDF Collaboration, T. Aaltonen et al., Phys. Rev. Lett. **100**, 082001 (2008).
 - [2] D0 Collaboration, V.M. Abazov et al., arXiv:0711.0319 [hep-ex].
 - [3] W.M. Yao et al., Particle Data Group, J. Phys. G **33**, 1 (2006).
 - [4] Z.G. Luo, X.L. Chen and X. Liu, arXiv:0901.0505 [hep-ph].
 - [5] W.A. Bardeen, E.J. Eichten and C.T. Hill, Phys. Rev. **D 68**, 054024 (2003); P. Colangelo, F. De Fazio and R. Ferrandes, Nucl. Phys. Proc. Suppl. **163**, 177 (2007).
 - [6] J. Lu, W.Z. Deng, X.L. Chen, and S.-L. Zhu, Phys. Rev. D **73**, 054012 (2006).
 - [7] The Babar Collaboration, B. Aubert et al., Phys. Rev. Lett. **90**, 242001 (2003); F. Porter, Eur. Phys. J. **C33**, 219 (2004).
 - [8] The Belle Collaboration, P. Krokovny et al., Phys. Rev. Lett. **91**, 262002 (2003).
 - [9] The CLEO Collaboration, D. Besson et al., Phys. Rev. **D 68**, 032002 (2003).
 - [10] S.L. Zhu, Int. J. Mod. Phys. **E 17**, 283-322 (2008).
 - [11] W.A. Bardeen, E.J. Eichten and C.T. Hill, Phys. Rev. **D 68**, 054024 (2003); M.A. Nowak, M. Rho and I. Zahed, Acta Phys. Polon. **B 35**, 2377 (2004); E. Kolomeitsev and M. Lutz, Phys. Lett. **B 582**, 39 (2004).
 - [12] K.T. Chao, Phys. Lett. **B 599**, 43 (2004).
 - [13] E. van Beveren and G. Rupp, Phys. Rev. Lett. **91**, 012003 (2003); E. van Beveren and G. Rupp, Eur. Phys. J. **C 32**, 493 (2004).
 - [14] Y. B. Dai, C. S. Huang, C. Liu, S. L. Zhu, Phys. Rev. D **68**, 114011 (2003).
 - [15] S. Narison, Phys. Lett. **B 605**, 319 (2005).
 - [16] Y. B. Dai, X. Q. Li, S. L. Zhu and Y. B. Zuo, Eur. Phys. J. C **55**, 249 (2008), arXiv: **hep-ph/0610327**.
 - [17] Y.Q. Chen and X.Q. Li, Phys. Rev. Lett. **93**, 232001 (2004); A.P. Szczepaniak, Phys. Lett. **B 567**, 23 (2003); T.E. Browder, S. Pakvasa and A.A. Petrov, Phys. Lett. **B 578**, 365 (2004).
 - [18] H.Y. Cheng and W.S. Hou, Phys. Lett. **B 566**, 193 (2003).
 - [19] T. Barnes, F.E. Close and H.J. Lipkin, Phys. Rev. **D 68** 054006 (2003); A.P. Szczepaniak, Phys. Lett. **B 567**, 23 (2003).
 - [20] A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli and A. D. Polosa, Phys. Rev. D **58**, 034004 (1998); A.D. Polosa, Riv. Nuovo Cim. **23N11**, 1, 2000.
 - [21] D. Ebert, T. Feldmann and H. Reinhardt, Phys. Lett. **B 388**, 154 (1996); T. Feldmann, arXiv: **hep-ph/9606451**; D. Ebert, T. Feldmann, R. Friedrich and H. Reinhardt, Nucl. Phys. **B 434** 619 (1995).
 - [22] A.F. Falk and M.E. Luke, Phys. Lett. **B 292**, 119 (1992).
 - [23] A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli and A.D. Polosa, Phys. Rev. **D 58**, 034004 (1998); A. Deandrea, R. Gatto, G. Nardulli and A.D. Polosa, Phys. Rev. **D 59**, 074012 (1999); A. Deandrea, R. Gatto, G. Nardulli and A.D. Polosa, J. High Energy Phys. **02**, 021 (1999); A. Deandrea, R. Gatto, G. Nardulli and A.D. Polosa, Phys. Rev. **D 61**, 017502 (1999); A.D. Polosa, arXiv: **hep-ex/9909371**.
 - [24] X.H. Guo, H.W. Ke, X.Q. Li, X. Liu and S.M. Zhao, Commun. Theor. Phys. **48**, 509-518 (2007), arXiv:hep-ph/0510146.
 - [25] X. Liu, Y.M. Yu, S.M. Zhao and X.Q. Li, Eur. Phys. J. **C 47**, 445 (2006), arXiv:hep-ph/0510146.
 - [26] S.M. Zhao, X. Liu and S.J. Li, Eur. Phys. J. **C 51**, 601-606 (2007), arXiv:hep-ph/0612008.
 - [27] A. Deandrea, G. Nardulli and A.D. Polosa, Phys. Rev. **D 68**, 097501 (2003).
 - [28] A. Deandrea, R. Gatto, G. Nardulli and A.D. Polosa, JHEP **9902 021** (1999).

- [29] M.Q. Huang, Phys. Rev. **D 69**, 114015 (2004); T.M. Aliev and M. Savci, Phys. Rev. **D 73** 114010 (2006); T.M. Aliev, K. Azizi and A. Ozpineci, Eur. Phys. J. **C 51**, 593-599 (2007), arXiv: [hep-ph/0608264](#).